

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <1/11>**  
**(excluding the port)**

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Schlegel's black rockfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2019	ND(4.0)	ND(3.2)	ND
Common skete (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2019	ND(3.7)	ND(3.6)	ND
Flatfish (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2019	ND(3.1)	ND(3.7)	ND
Chub mackerel (muscle)	Around 1km Offshore of Ota River (T-S1)	May 17, 2019	ND(3.8)	ND(3.1)	ND
Blue crab (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.5)	ND(4.1)	ND
Lepidotrigla microptena (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.1)	ND(3.5)	ND
Yellow gosefish (whole)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.8)	ND(3.1)	ND
Schlegel's black rockfish (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(2.7)	ND(3.8)	ND
Common skete (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.7)	ND(2.9)	ND
Microstomus achne (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.7)	ND(4.0)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <2/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish ① (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(2.5)	ND(3.6)	ND
Flatfish ② (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.5)	ND(3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(4.0)	ND(4.1)	ND
Chub mackerel (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.4)	ND(3.0)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Odaka Ward (T-S2)	May 17, 2019	ND(3.6)	ND(3.1)	ND
Spiny dogfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(3.1)	ND(3.8)	ND
Stone flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(4.0)	ND(4.1)	ND
Yellow goosefish (whole)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(4.2)	ND(3.4)	ND
Common skete (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(3.2)	ND(4.3)	ND
Flatfish (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(4.2)	ND(3.5)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

**Nuclide Analysis Results of Fish and Shellfish**  
**(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <3/11>**  
**(excluding the port)**

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Searobin (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(2.8)	ND(3.2)	ND
Marbled sole (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(3.5)	ND(3.5)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Ukedo River (T-S3)	May 10, 2019	ND(3.7)	ND(3.6)	ND
Stone flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(4.1)	ND(4.3)	ND
Japanese angel shark (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(2.9)	3.9	3.9
Lepidotrigla microptena (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.5)	ND(3.3)	ND
Yellow goosefish (whole)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(4.5)	ND(4.1)	ND
Common skate (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(4.3)	ND(3.5)	ND
Microstomus achne (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.5)	3.5	3.5
Flatfish ① (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.0)	ND(3.3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <4/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish ② (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.6)	ND(3.7)	ND
Searobin (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.7)	ND(3.3)	ND
Littlemouth flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(4.1)	ND(4.1)	ND
Roundnose flounder (muscle)	Around 3km Offshore of Fukushima Daiichi (T-S4)	May 10, 2019	ND(3.9)	ND(3.7)	ND
Greenling (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(4.3)	ND(3.7)	ND
Spiny dogfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.8)	ND(4.0)	ND
Stone flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(4.0)	ND(3.8)	ND
Common skete (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.8)	ND(3.5)	ND
Microstomus achne (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.5)	ND(3.1)	ND
Flatfish (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.6)	ND(3.7)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <5/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Marbled sole (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.7)	ND(3.9)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Kido River (T-S5)	May 28, 2019	ND(3.9)	ND(3.6)	ND
Greenling (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	ND(3.5)	5.1	5.1
Spiny dogfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	ND(3.9)	ND(3.7)	ND
Schlegel's black rockfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	6.7	95	101.7 ※ <sup>1</sup>
Microstomus achne (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	ND(3.8)	ND(3.1)	ND
Flatfish (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	ND(3.4)	ND(3.7)	ND
Roundnose flounder (muscle)	Around 2km Offshore of Fukushima Daini (T-S7)	May 28, 2019	ND(4.4)	ND(3.5)	ND
Stone flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(4.4)	ND(3.6)	ND
Yellow gosefish (whole)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.9)	ND(3.6)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Analyzed by: Tokyo Power Technology Ltd.

※<sup>1</sup> Measured values are compared with the baseline to two significant figures. (Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.) The measurement result this time is 100Bq/kg as rounded to two significant digits, and does not exceed the baseline. This has been announced in June 14,2019.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <6/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.8)	ND(3.5)	ND
Microstomus achne (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.9)	4.9	4.9
Flatfish ① (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.0)	ND(3.9)	ND
Flatfish ② (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.3)	ND(3.9)	ND
Marbled sole (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.5)	ND(4.0)	ND
Roundnose flounder (muscle)	Around 4km Offshore of Kumagawa (T-S8)	May 28, 2019	ND(3.9)	ND(3.3)	ND
Greenling (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.7)	ND(3.3)	ND
Stone flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.7)	ND(3.9)	ND
Yellow goosfish (whole)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.9)	ND(4.3)	ND
Microstomus achne (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(4.5)	ND(3.6)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <7/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Flatfish (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(4.5)	ND(3.7)	ND
Common Japanese conger (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(4.5)	ND(4.1)	ND
Littlemouth flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.8)	ND(3.9)	ND
Pacific cod (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.1)	ND(4.0)	ND
Ridged-eye flounder (muscle)	Around 15km Offshore of Odaka Ward (T-B1)	May 14, 2019	ND(3.9)	ND(3.4)	ND
Spiny dogfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(4.0)	ND(3.4)	ND
Yellow goosfish (whole)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.2)	ND(3.5)	ND
Common skate (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(4.3)	ND(3.5)	ND
Microstomus achne (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.6)	ND(3.9)	ND
Flatfish (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(4.0)	ND(3.8)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <8/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Littlemouth flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(2.8)	ND(3.7)	ND
Marbled sole (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.2)	ND(3.7)	ND
Chub mackerel (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.6)	ND(3.4)	ND
Pacific cod (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.5)	ND(3.8)	ND
Roundnose flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.8)	ND(3.1)	ND
Ridged-eye flounder (muscle)	Around 18km Offshore of Ukedo River (T-B2)	May 14, 2019	ND(3.7)	ND(4.3)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.9)	ND(3.7)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.4)	ND(3.5)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.6)	ND(3.4)	ND
Yellow goosefish (whole)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.2)	ND(4.2)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.



Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <9/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Common skate (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.3)	ND(3.5)	ND
White croaker (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.4)	ND(3.7)	ND
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.1)	ND(2.9)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(4.2)	ND(3.7)	ND
Flatfish ① (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.4)	ND(3.7)	ND
Flatfish ② (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.2)	ND(3.2)	ND
Searobin (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.9)	ND(3.9)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.4)	ND(3.5)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.3)	ND(3.6)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(4.5)	ND(3.3)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <10/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.1)	ND(3.4)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daiichi (T-B3)	May 7, 2019	ND(3.5)	ND(4.7)	ND
Greenling (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.6)	ND(3.9)	ND
Stone flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.3)	ND(4.0)	ND
Lepidotrigla microptena (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.6)	ND(3.0)	ND
Yellow goosfish (whole)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.2)	ND(4.5)	ND
Common skate (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.3)	ND(4.3)	ND
White croaker (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.6)	ND(3.6)	ND
Pointhead flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.0)	ND(3.8)	ND
Cloudy catshark (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.6)	ND(4.1)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.

Nuclide Analysis Results of Fish and Shellfish  
(Ocean Area Within 20km Radius of Fukushima Daiichi Nuclear Power Station) <11/11>  
(excluding the port)

Name of Sample (Region)	Place of Sampling (Place No.)	Date of Sampling	Radioactivity Concentration [Bq/kg (Raw)] (Half-life)		
			Cs-134 (Approx. 2 years)	Cs-137 (Approx. 30 years)	CS (Sum)
Drumfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.3)	ND(3.1)	ND
Microstomus achne (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.9)	ND(3.6)	ND
Flatfish (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.1)	ND(4.2)	ND
Common Japanese conger (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.7)	ND(3.9)	ND
Littlemouth flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(4.4)	ND(4.1)	ND
Marbled sole (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.9)	ND(4.1)	ND
Roundnose flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.4)	ND(4.0)	ND
Ridged-eye flounder (muscle)	Around 10km Offshore of Fukushima Daini (T-B4)	May 7, 2019	ND(3.9)	ND(3.6)	ND

\*When analyzed results are less than detection limits of radioactivity concentrations, the values are showed as "ND." Detection limits of individual nuclides are shown in parenthesis.

\*Since April 1, 2012, the baseline is the sum of radioactivity concentrations of Cs-134 and Cs-137, which is 100Bq/kg.

\*Analyzed by: Tokyo Power Technology Ltd.